

NEWS

Music evokes emotion in children with autism

BY ANDREA ANDERSON

23 MAY 2008

Music to the ears: Imaging studies may explain why children with autism respond to emotions conveyed through music better than to other forms of communication.

Children with autism struggle to understand social and emotional cues from other people's actions or words: that is one of autism's cardinal features, in fact.

These same children respond to music, however, understanding emotions conveyed through non-verbal musical cues. And music therapy has been shown to improve symptoms of autism such as verbal communication, agitation and social interaction deficits¹.

Some children with autism also have exceptional musical abilities, such as pitch sensitivity².

It can be difficult to tease apart which parts of the brain process emotion conveyed by music, and the biology underlying this apparent anomaly in children with autism has been a mystery.

Scientists are turning to brain imaging to understand why these children understand music better than they do other forms of social communication.

"If they can do it for music, why can't they do it for faces? For voices?" asks **Istvan Molnar-**

Szakacs, a research neuroscientist at the University of California, Los Angeles (UCLA) Tennenbaum Center for the Biology of Creativity.

"I feel like what the autistic children are really getting out of music is a salient emotional charge that is temporally organized and structured, so it comes in a form that is relevant for them," he says.

In an upcoming study funded by a \$40,000 grant from the **GRAMMY Foundation**, Molnar-Szakacs is planning functional magnetic resonance imaging (fMRI) studies on 15 high-functioning children with autism between the ages of 9 and 13, and 15 age- and sex-matched typically developing controls.

Emotional excerpts:

A coronal view of the brain using fMRI, illustrating activity within the mirror neuron system, thought to be essential for emotional response to music.

Musical colors: A coronal view of the brain using fMRI, illustrating activity within the mirror neuron system, thought to be essential for emotional response to music.

The researchers will ask each child to identify emotions from pictures of faces or from 12- to 20-second emotional musical excerpts ? long enough to convey emotion, but not so long that children will move in the scanner.

Though there is a six-second delay between the brain response and the time the signal is measured, the fMRI effectively provides real-time information about areas of the brain responding to music.

The children will also be asked to provide feedback based on what they've heard or seen ? for instance, pushing a button to indicate whether the face or the piece of music is happy or sad.

The researchers are still deciding on musical excerpts, Molnar-Szakacs says, but the pieces are

likely to be compositions without lyrics that could confound the results.

The project is expected to begin this fall after pilot behavioral experiments tease out and optimize the protocol. "At this point we're just trying to figure out what the best set of stimuli are that children resonate with the most," he says.

Molnar-Szakacs and his colleagues are particularly interested in exploring links between music and two brain networks: the limbic system and the mirror neuron system.

The limbic system is an interconnected series of brain structures ? including the hippocampus and amygdala ? that process emotions such as happiness and fear. Mirror neurons are cells in the premotor cortex and parietal cortex that fire when people watch or imitate other people?s actions and interpret their intentions.

Although mirror neurons are most often associated with seeing and understanding actions, there is evidence that they may also help decipher meaning and emotion in music³.

Musical movements:

Evolutionarily, Molnar-Szakacs says, making music is intimately linked to movement ? from drumming to tapping toes and performing ceremonial dances. Recorded music has turned it into a more passive process.

"Before MP3 players, people had to move their arm or leg or vocal apparatus to generate music," he says. "But the brain still uses systems that have evolved to process action to understand music."

That may not necessarily be the case in children with autism. The potential link to mirror neurons is intriguing because several studies have proposed that children with autism have decreased or altered mirror neuron activity^{4,5}.

The research may help understand whether children with autism respond to the same auditory cues that produce emotion in others, notes **Vinod Menon**, a researcher in psychiatry and neuroscience at Stanford University. "Certainly one could make very strong predictions" that brain areas would respond differently to music in those with autism, he says.

Still, mirror neurons could respond to many different elements of musical stimuli, Menon cautions. It may also be difficult to tease apart the function of mirror neurons from other brain regions that respond to music.

Because people listen to music for its emotive force, it is also important to explore reward and pleasure systems areas of the brain, he says.

For example, Menon's research indicates that listening to pleasant music affects brain areas called the nucleus accumbens and the ventral tegmental area – regions involved in reward processing⁶.

Previous studies on music and children with autism have been done on the behavioral level, so there is little information to help tease apart the specific roles of overlapping brain systems.

If reward centers are involved, for example, "that's something that would come out of the imaging," Molnar-Szakacs says. "We're just getting started."

References:

1.

Boso M. *et al. J. Altern. Complement. Med.* **13**, 709-712 (2007) [PubMed](#)

2.

Heaton P.J. *Autism Dev. Disord.* **35**, 787-793 (2005) [PubMed](#)

3.

Molnar-Szakacs I. and K. Overy *SCAN* **1**, 235-241 (2006) [Abstract](#)

4.

Dapretto M. *et al. Nat. Neurosci.* **9**, 28-30 (2006) [PubMed](#)

5.

Cattaneo L. *et al. Proc. Natl. Acad. Sci. USA* **104**, 17825-17830 (2007) [PubMed](#)

6.

Menon V. and D.J. Levitin *Neuroimage* **28**, 175-184 (2005) [PubMed](#)

